THERE ARE AROUND

ADULTS WHO SMOKE CIGARETTES IN GREAT BRITAIN.

THAT'S ALMOST 17% OF THE ENTIRE UK POPULATION.

ABOUT HALF OF ALL REGULAR **SMOKERS** WILL EVENTUALLY BE KILLED BY THEIR ADDICTION.

SMOKING COSTS THE NHS APPROX **£2.7BN** A YEAR FOR TREATING DISEASES CAUSED BY SMOKING.

IN 2013, UK SMOKERS SPENT AROUND £14BN ON TOBACCO.



**£2.7**BN

£14BN

**TOBACCO** 

CIGARETTES ARE THE MAIN CAUSE OF FATAL ACCIDENTAL FIRES IN THE HOME.

IN 2008, SMOKERS' MATERIALS ACCOUNTED FOR



ION-FATAL CASUALTIES FROM FIRES IN THE HOME.

**PERCENTAGE OF ADULT** SMOKERS.

	22%	MEN
	19%	WOMEN

**PERCENTAGE** OF EX-SMOKERS.



**ABOUT TWO-**THIRDS OF



CURRENT SMOKERS WOULD LIKE TO STOP SMOKING.

ON AVERAGE, CIGARETTE SMOKERS DIE 10 YEARS YOUNGER THAN NON-SMOKERS.





SMOKING AFFECTS YOUR SENSE OF TASTE AND SMELL SMOKERS ARE MORE LIKELY TO DEVELOP WRINKLES YOUNGER AND HAVE **DENTAL PROBLEMS**.

A 20-A-DAY SMOKER WILL SPEND AROUND

SMOKERS UNDER THE AGE OF 40 HAVE A FIVE **TIMES GREATER RISK** OF A HEART ATTACK THAN NON-SMOKERS.



SMOKING CAUSES AROUND OF DEATHS FROM UNG CANCER AND AROUND **5U** 70 OF **DEATHS FROM BRONCHITIS** AND EMPHYSEMA.

IF SMOKERS QUIT BEFORE THE AGE OF 30, THEY CAN AVOID ALMOST ALL OF THE RISK OF LUNG CANCER ATTRIBUTABLE TO SMOKING.



EVERY YEAR, OVER 100,000 **SMOKERS** IN THE UK DIE FROM **SMOKING** RELATED CAUSES.



Source: Action for Smoking and Health (www.ash.org.uk)